arXiv:1108.0830v2 [math.NT] 24 Dec 2011

EMBEDDINGS OF FIELDS IN SIMPLE ALGEBRAS OVER GLOBAL FIELDS

SHENG-CHI SHIH, TSE-CHUNG YANG AND CHIA-FU YU

Abstract. Let F be a global field, A a central simple algebra over F and K a finite (separable or not) field extension of F with degree $[K:F]$ dividing the degree of A over F. An embedding of K in A over F exists implies an embedding exists locally everywhere. In this paper we give a quite complete account under what conditions the converse (i.e. the local-global principle in question) may hold.

1. Introduction

The topic on central simple algebras over global fields is one of central themes in number theory. One way to extract information of a central simple algebra is to study its maximal subfields. For example, this information would be useful when one attempts to analyze the terms in the geometric side of trace formulas. Indeed, in the geometric side of the trace formula for the multiplicative group of a given central simple algebra, certain major terms (elliptic ones) are described by conjugacy classes of these maximal subfields. This leads to a very important approach, due to Eichler [3], to calculating the class number (of an open compact level subgroup), as the class number can be written as the trace of the characteristic function supported on this level subgroup. In this pioneer work [3] Eichler established results in the trace formula for computing the class numbers and the type numbers of a maximal order or a hereditary order in a given quaternion algebra over any global field. He reduced some of the problem of computing class numbers to computing so called the *optimal embeddings* among orders in maximal subfields and in the quaternion algebras, which are of purely local nature. See Eichler [3] and Vignéras [9] for more details.

A useful tool to study maximal subfields is the Hasse principle. This enables us to describe the properties of these subfields from local information. A naive attempt along the same direction is whether one can describe subfields of a central simple algebra over a global field F from local information. In this paper we study embeddings a finite field extension of F, which does not necessarily have the maximal degree, in a central simple algebra over F, and attempted to clarify related problems in this situation. More precisely, consider a finite-dimensional central simple algebra A over a global field F and a finite (separable or not) field extension K of F whose degree divides the degree of A over K, that is, $[K:F] \mid \sqrt{[A:F]}$. Naturally one considers the following basic questions:

(Q1) What is the necessary and sufficient condition for which the field K can be embedded into A over F?

Date: December 30, 2011.

(Q2) What is the necessary and sufficient condition for which the F_v-algebra K_v can be embedded into the F_v-algebra A_v over F_v? Here v is a place of F, F_v denotes the completion of F at v, $K_v := K \otimes_F F_v$ and $A_v := A \otimes_F F_v$.

(Q3) Does the local-global principle for embedding the field K into A as F-algebras hold? That is, if K_v can be embedded in A_v for all places v of F, then can K be embedded in A?

We say that the Hasse principle for a pair (K, A) as above holds if the question (Q3) for K and A has a positive answer. When the field extension K has the maximal degree $\deg A$, this result is well-known and useful; see Pierce [4, Section 18.4], Prasad-Rapinchuk [5, Proposition A.1] and [6, Proposition 2.7]. However, the case when K does not have the maximal degree is not explored in the literature.

We now outline the contents of this paper. In the first part of this paper we answer the questions (Q1), (Q2) and (Q3). In [11], the third author of the present paper studies the problem of embeddings of one semi-simple algebra into another one over an arbitrary ground field. In Section 2 we recall some results of embeddings obtained in [11] and provide some proofs of them for the reader's convenience. These give rise to a numerical criterion for the question (Q1) over any field; see Lemma 2.4 for the precise statement. In Section 3 we apply this embedding result to the case when the base field is either a local field or a global field. This yields a more explicit numerical criterion for (Q1) and (Q2), respectively. Using these criteria, we then show that there are a Galois extension K over F of degree 8 and a central simple algebra A over F of degree 24 for which the Hasse principle does not hold; see an example in Section 3.4. Also see Proposition 1.4 below for other cases for which the Hasse principle fails .

We now describe the results in this part. Let K and A be as above. Let $A = \mathrm{Mat}_n(\Delta)$, where Δ is the division part of A and $n := \mathrm{c}(A)$ is called the capacity of A (see Definition 2.1). For a place v of F, let $K_v := K \otimes_F F_v = \prod_{w|v} K_w$ be the product of local fields. Let V^F and V^K denote the sets of places of F and K, respectively. Let d_v be the order of the class $[A_v]$ in the Brauer group of F_v.

Theorem 1.1.

(1) *The set of $A_v^\times$-conjugacy classes of F_v-embeddings of K_v in A_v is in bijection with the following set*

$$\mathcal{E}_{F_v}(K_v, A_v) := \left\{ (x_w)_{w|v} \mid x_w \in \mathbb{N},\ \sum_{w|v} \ell_w x_w = \mathrm{c}(A_v) \right\},$$

where

$$\ell_w := [K_w : F_v] / \gcd([K_w : F_v], \deg(\Delta)).$$

In particular, there is an F_v-embedding of K_v in A_v if and only if the finite set $\mathcal{E}_{F_v}(K_v, A_v)$ is non-empty.

(2) *There is an F-embedding of K in A if and only if*

$$[K : F] \mid n\, \mathrm{c}(\Delta \otimes_F K),$$

where $\mathrm{c}(\Delta \otimes_F K)$ is the capacity of the central simple algebra $\Delta \otimes_F K$ over K.

(3) *The number $\mathrm{c}(\Delta \otimes_F K)$ can be computed as follows.*

  (i) *The degree δ_0 of Δ is the least common multiple of d_v for all $v \in V^F$.*

(ii) *For $w|v$, the order d'_w of the class $[A_v \otimes_{F_v} K_w]$ in the Brauer group of K_w is given by*

$$d'_w = d_v/\gcd([K_w : F_v], d_v).$$

(iii) *The order δ' of the class $[A \otimes_F K]$ in the Brauer group of K is the least common multiple of d'_w for all $w \in V^K$.*

(iv) *The capacity $\mathrm{c}(\Delta \otimes_F K)$ is then given by δ_0/δ'.*

The proof of Theorem 1.1 is given in Sections 3.1 and 3.2; see especially Propositions 3.2, 3.3 and 3.4.

In the last part of Section 3, we give a necessary and sufficient condition for a pair (K, A) so that the Hasse principle in question holds. We associate to each pair (K, A) an element

$$\bar{\mathbf{x}} = (\bar{\mathbf{x}}_w)_{w \in V^K} \in \bigoplus_{w \in V^K} \mathbb{Q}/\mathbb{Z}$$

as follows. For any $w \in V^K$, put

$$\mathbf{x}_w := \frac{\mathrm{c}(A_v) \cdot \gcd([K_w : F_v], d_v)}{[K : F]} \in \mathbb{Q}_{>0}$$

and let $\bar{\mathbf{x}}_w$ be the class of $\mathbf{x}_w$ in $\mathbb{Q}/\mathbb{Z}$, where v is the place of F below w.

We show that this element $\bar{\mathbf{x}}$ is an obstruction class to the Hasse principle for (K, A), i.e. if this class does not vanish, then the Hasse principle for (K, A) fails. We also show that the vanishing of $\bar{\mathbf{x}}$ is the only obstruction to the Hasse principle. Namely, we have the following result (Theorem 3.6).

Theorem 1.2. *Notations as above. An F-embedding of K in A exists if and only if an F_v-embedding of K_v in A_v exists for all $v \in V^F$ and that the element $\bar{\mathbf{x}}$ vanishes.*

The second part of this paper deals with the Hasse principle for a family of pairs (K, A) with both degrees $[K : F]$ and $\deg(A)$ constant. Let (k, δ) be a pair of positive integers with $k \mid \delta$. We say that the Hasse principle for the pair (k, δ) holds if for any finite field extension K over F of degree k and any central simple algebra A over F of degree δ, the Hasse principle for the pair (K, A) holds.

We describe the results in this part. Let (k, δ) be a pair of positive integers as above. For each partition λ of k

$$\lambda = (k_1, \dots, k_t), \quad k_1 \le \dots \le k_t, \quad \sum_{i=1}^{t} k_i = k$$

and a pair $\mu = (s, d)$ of two positive integers s and d with $\delta = sd$, we definite a finite set

$$\mathcal{E}(\lambda, \mu) := \{(x_i) \in \mathbb{N}^t \mid \sum_{i=1}^{t} \ell_i x_i = s\},$$

where $\ell_i := k_i/(k_i, d)$, and a vector $\mathbf{x}(\lambda, \mu)$ in $\mathbb{Q}^t$

$$\mathbf{x}(\lambda, \mu) := (\mathbf{x}_i), \quad \mathbf{x}_i = \mathbf{x}_i(\lambda, \mu) := s(k_i, d)/k, \quad \forall\, i = 1, \dots, t.$$

Here (k_i, d) denotes the greatest common divisor of the integers k_i and d; to indicate instead a pair of integers, say k and δ, we write *the pair* (k, δ).

Let $\mathrm{LD}(k, \delta)$ be the set consisting of all pairs (λ, μ), where

- λ is a partition of k, and

- $\mu = (s, d)$ is a decomposition of δ into the product of two positive integers,

such that the set $\mathcal{E}(\lambda, \mu)$ is non-empty. ("LD" stands for all possible local decompositions). The following result gives a simple way to check the Hasse principle for a pair (k, δ).

Theorem 1.3. *Let (k, δ) be a pair of positive integers with $k \mid \delta$.*

1. *Suppose for all elements $(\lambda, \mu) \in \mathrm{LD}(k, \delta)$, the vector $\mathbf{x}(\lambda, \mu)$ has integral components. Then the Hasse principle for (k, δ) holds.*
2. *Suppose there is an element $(\lambda, \mu) \in \mathrm{LD}(k, \delta)$ such that $\mathbf{x}_i(\lambda, \mu) \notin \mathbb{N}$ for some i. Then there are a finite separable field extension K of degree k over F and a central division algebra A of degree δ over F so that the Hasse principle for (K, A) fails. In particular, the Hasse principle for the pair (k, δ) fails.*

The proof of Theorem 1.3 uses Theorem 1.2, the Hilbert irreducibility theorem and the global class field theory. Based on Theorem 1.3, we prove the following result (Proposition 4.6 and Corollary 4.7), which also shows that (Q3) has a negative answer in general.

Proposition 1.4.

1. *Let $\delta = p_1^{n_1} \cdots p_r^{n_r}$, and let k be a positive integer with $k > 1$ and $k \mid \delta$, where each p_i is a prime number. Assume that $k \le \delta / p_i^{n_i}$ and $p_i \mid k$ for some $1 \le i \le r$ (so $r \ge 2$). Then the Hasse principle for the pair (k, δ) does not hold.*
2. *Let δ be a positive integer divisible by two primes. Then there is a positive integer k with $k \mid \delta$ such that the Hasse principle for the pair (k, δ) does not hold.*

Clearly Proposition 1.4 (2) follows from Proposition 1.4 (1). Proposition 1.4 tells us that when δ is divisible by two primes and k is "small" comparable to δ, the Hasse principle for (δ, k) fails. It is also well-known that when $k = \delta$, the maximal degree case, the Hasse principle holds. Therefore, it is natural to ask the Hasse principle when δ is a prime power or k is "very close to δ". So we consider the following two more questions:

(Q4) Does the Hasse principle for a pair (k, δ) hold if $\delta = p^n$ is a prime power?
(Q5) Does the Hasse principle for a pair (k, δ) hold if δ / k is a prime?

About the question (Q4), we prove the following result (Theorem 5.4).

Theorem 1.5. *Let $(k, \delta) = (p^m, p^n)$ be a pair of powers of a prime number p. The Hasse principle for the pair $(k, \delta) = (p^m, p^n)$ holds if and only if one of the following conditions holds:*

(i) *$m = n$.*
(ii) *$m = 1$ or 2.*

About the question (Q5), we first prove the following (Proposition 5.8).

Proposition 1.6. *Let (k, δ) be a pair of positive integers such that $p_1 = \delta / k$ is a prime. Suppose that p_1 is not the smallest prime divisor of δ. Then the Hasse principle for the pair (k, δ) does not hold.*

For the remaining case where p_1 is the smallest prime divisor, the situation is more complicated. Therefore, we restrict ourselves to the case where δ is divisible by exact two primes, i.e. $\delta = p_1^{n_1} p_2^{n_2}$, where p_1 and p_2 are primes. In this case, we are able to give the complete answer. We prove (Theorem 5.14)

Theorem 1.7. *Let* $(\delta, k) = (p_1^{n_1} p_2^{n_2}, p_1^{n_1-1} p_2^{n_2})$*, where* $p_1 < p_2$ *are primes and* $n_1, n_2 \in \mathbb{N}$*. The Hasse principle for the pair* (k, δ) *holds if and only if* $(n_1, n_2) = (1, 1)$*.*

The proof of Theorem 1.7 is divided into 5 different cases and we prove them case by case (from Propositions 5.9 to 5.13).

The following result (Theorem 5.15) is a consequence of Proposition 1.4 and Theorem 1.5.

Theorem 1.8. *The positive integers* δ *with the property that for any central simple algebra* A *over* F *of degree* δ *and any finite field extension* K *of degree* $[K : F]$ *dividing* δ *the Hasse principle for* (K, A) *holds are* $1, p, p^2, p^3$*, where* p *is a prime.*

In the last part of this paper, we study the embedding problem from the topological point of view. Assume that K is *separable.* Let $\mathfrak{F}$ be a separable closed field which contains the fields F_v for all places $v \in V^F$; such a big field $\mathfrak{F}$ exists; see Section 6.

Suppose we are given an F_v-embedding $\iota_v : K_v \hookrightarrow A_v$ for each place $v \in V^F$ with the following compatibility condition. Let

$$\iota_v \otimes \mathrm{id}_{\mathfrak{F}} : K_v \otimes_{F_v} \mathfrak{F} \hookrightarrow A_v \otimes_{F_v} \mathfrak{F}$$

be the induced $\mathfrak{F}$-embedding. Noting that $K_v \otimes_{F_v} \mathfrak{F} = K \otimes_F \mathfrak{F}$ and $A_v \otimes_{F_v} \mathfrak{F} = A \otimes_F \mathfrak{F}$, we have

$$\iota_v \otimes \mathrm{id}_{\mathfrak{F}} : K \otimes_F \mathfrak{F} \hookrightarrow A \otimes_F \mathfrak{F}.$$

The compatibility condition requires that for any two places $v_1, v_2 \in V^F$, the induced maps $\iota_{v_1} \otimes \mathrm{id}_{\mathfrak{F}}$ and $\iota_{v_2} \otimes \mathrm{id}_{\mathfrak{F}}$ are conjugate by an element in $(A \otimes_F \mathfrak{F})^\times$. The question is: is there an F-embedding $\iota : K \hookrightarrow A$ such that the induced map $\iota \otimes \mathrm{id}_{F_v}$ is conjugate to ι_v for all $v \in V^F$? We show this question has an affirmative answer (Theorem 6.7).

Theorem 1.9. *Let* $\iota_v : K_v \hookrightarrow A_v$ *be an* F_v*-embedding of* K_v *into* A_v *for each* $v \in V^F$*. Suppose that for any two places* $v_1, v_2 \in V^F$*, the maps* $\iota_{v_1} \otimes \mathrm{id}_{\mathfrak{F}}$ *and* $\iota_{v_2} \otimes \mathrm{id}_{\mathfrak{F}}$ *are conjugate in* $(A \otimes_F \mathfrak{F})^\times$*. Then there exist an* F*-embedding* $\iota : K \hookrightarrow A$ *such that* $\iota \otimes \mathrm{id}_{F_v}$ *is conjugate to* ι_v *in* $A_v^\times$ *for all* $v \in V^F$*.*

The proof of Theorem 1.9 uses Theorems 1.1, 1.2 and the Chebotarev density theorem.

We hope this paper may provide some useful information to the geometric side of the trace formula mentioned above. We also remark that the local-global principle for embedding of maximal subfields in central simple algebras over global fields *with involutions*, which is more involved, is studied in Prasad and Rapinchuk [6].

This paper is organized as follows. In Section 2 we collect and show some general embedding results over any field based on [11]. In Section 3, we give a more detailed study about embeddings of a field extension K into a central simple algebra A over a global field. We determine for which pair (K, A) the Hasse principle holds. In particular, we answer the questions (Q1), (Q2) and (Q3). In Sections 4 and 5, we work on the Hasse principle for a given pair (k, δ) of positive integers with k dividing δ. In Section 6, we give the proof of Theorem 1.9.

2. GENERAL EMBEDDING RESULTS

2.1. **Setting.** Let F denote the ground field, which is arbitrary in this section. All F-algebras considered in this paper are assumed to be finite-dimensional as F-vector spaces and have the identity. As the standard convention, an F-algebra homomorphism is a ring homomorphism over F; in particular, it sends the identity of the source to the identity of the target.

We recall the following definition for central simple algebras; see [7].

Definition 2.1. The *degree*, *capacity*, and *index* of a central simple algebra A over F are defined as

$$\deg(A) := \sqrt{[A:F]}, \quad \mathrm{c}(A) := n, \quad \mathrm{i}(A) := \sqrt{[\Delta : F]},$$

if $A \cong \mathrm{Mat}_n(\Delta)$, where Δ is a division algebra over F, which is uniquely determined by A up to isomorphism. The algebra Δ is also called the *division part* of A.

For the convenience of discussion, we introduce the following definition (cf. [11, Definition 2.1]).

Definition 2.2.

(1) Let V be a finite-dimensional vector space over F, and A an (finite-dimensional) arbitrary F-algebra. We say that V *is A-modulable* if there is a right (or left) A-module structure on V. If B is any F-subalgebra of A and V is already a right (resp. left) B-module, then by saying V is A-modulable we mean that the right (resp. left) A-module structure on V is required to be compatible with the underlying B-module structure on V.

(2) For any two F-algebras A_1, A_2, let $\mathrm{Hom}_F(A_1, A_2)$ denote the set of F-algebra homomorphisms from A_1 to A_2, and let

$$\mathrm{Hom}_F^*(A_1, A_2) \subset \mathrm{Hom}_F(A_1, A_2)$$

be the subset consisting of embeddings of A_1 in A_2. For two maps $\varphi_1, \varphi_2 \in \mathrm{Hom}_F(A_1, A_2)$, we say φ_1 and φ_2 are *equivalent* if there is an element $b \in A_2^\times$ such that $\varphi_2 = \mathrm{Int}(b) \circ \varphi_1$. That is, $\varphi_2(a) = b\,\varphi_1(a)\,b^{-1}$ for all $a \in A_1$. Then $A_2^\times \backslash \mathrm{Hom}_F(A_1, A_2)$ is the set of equivalence classes of F-algebra homomorphisms from A_1 to A_2. Write

$$\mathcal{O}_{A_1,A_2} := A_2^\times \backslash \mathrm{Hom}_F(A_1, A_2) \quad \text{and} \quad \mathcal{O}^*_{A_1,A_2} := A_2^\times \backslash \mathrm{Hom}^*_F(A_1, A_2)$$

for the orbits spaces.

We often simply write $A_1 \otimes A_2$ for $A_1 \otimes_F A_2$ if the ground field F is understood.

2.2. **General embedding lemmas.** Let A be a central simple algebra over F. We realize $A = \mathrm{End}_\Delta(V)$, where Δ is the division part of A and V is a right Δ-module. Let A' be another simple F-algebra , not necessarily central over F, with center K. Then there is an F-algebra homomorphism $\varphi : A' \to A$ if and only if there is an

(A', Δ)-bimodule structure on V, or V is $\Delta \otimes_F A'^o$-modulable. Here A'^o denotes the opposite algebra of A'. Suppose $n := \dim_\Delta V$ and

$$\Delta \otimes_F A'^o \simeq (\Delta \otimes_F K) \otimes_K A'^o \simeq \mathrm{Mat}_c(\Delta'),$$

where Δ' is the division part of this central simple algebra over K. We have

$$[\Delta : F][A' : F] = c^2[\Delta' : F], \quad \text{and} \quad \dim_F V = n[\Delta : F]. \tag{2.1}$$

Then V is $\Delta \otimes_F A'^o$-modulable if and only if

$$c[\Delta' : F] \mid \dim_F V. \tag{2.2}$$

This is equivalent to $[\Delta : F][A' : F] \mid cn[\Delta : F]$, or equivalently

$$[A' : F] \mid nc. \tag{2.3}$$

We have proved

Lemma 2.3. *Let A be a central simple F-algebra and A' a simple F-algebra. Suppose $A \simeq \mathrm{Mat}_n(\Delta)$. Then there exists an embedding of A' in A if and only if*

$$[A' : F] \mid n \cdot \mathrm{c}(\Delta \otimes_F A'^o). \tag{2.4}$$

Next we consider the case where A' is semi-simple instead. Write $A' = \prod_{i=1}^s A'_i$ into simple factors and let K_i be the center of A'_i. The existence of an embedding of A' in A is equivalent to that there is an (A', Δ)-bimodule structure on V. This means that there is a decomposition of V into Δ-submodules

$$V = V_1 \oplus \cdots \oplus V_s \tag{2.5}$$

so that each V_i is a *non-zero* (A'_i, Δ)-bimodule. Put $n_i := \dim_\Delta V_i$ and let c_i be the capacity of the central simple algebra

$$\Delta \otimes_F A'^o_i = (\Delta \otimes_F K_i) \otimes_{K_i} A'^o_i$$

over K_i. Then we get the conditions

$$n = \sum_{i=1}^s n_i, \quad \text{and} \quad [A'_i : F] \mid n_i c_i, \quad \forall\, i = 1, \dots, s. \tag{2.6}$$

Lemma 2.4. *Let A and $A' = \prod_{i=1}^s A'_i$ be as above. Then there is an embedding of the F-algebra A' into A if and only if there are positive integers n_i for $i = 1, \dots, s$ such that*

$$n = \sum_{i=1}^s n_i, \quad \textit{and} \quad [A'_i : F] \mid n_i c_i, \quad \forall\, i = 1, \dots, s,$$

where c_i the capacity of the central simple algebra $\Delta \otimes_F A'^o_i$ over K_i.

Lemma 2.5. *Let A and $A' = \prod_{i=1}^s A'_i$ be as above. Let φ, φ' be two maps in $\mathrm{Hom}_F(A', A)$, and let V_φ and $V_{\varphi'}$ be the induced (A', Δ)-bimodules underlying the space V. Then φ and φ' are equivalent if and only if the (A', Δ)-bimodules V_φ and $V_{\varphi'}$ are isomorphic.*

PROOF. See [11, Lemma 3.2]

2.3. **Maximal degree field extension case.** We apply Lemmas 2.3 and 2.4 to the case where the semi-simple algebra A' is commutative and obtain the following well-known result. This is also a consequence of a result of Chuard-Koulmann and Morales [2, Proposition 4.3].

Lemma 2.6.

(1) *Let A be a central simple algebra over F and K is a field extension of F with $[K:F]=\deg(A)$. Then there exists an embedding of K into A if and only if K splits A.*

(2) *Let A be a central simple algebra over F and $K=\prod_{i=1}^{s} K_i$ is commutative semi-simple F-algebra with $[K:F]=\deg(A)$. Then there exists an embedding of K into A if and only if each K_i splits A.*

Proof. (1) Be Lemma 2.3, the set $\mathrm{Hom}_F(K,A)$ is non-empty if and only if $[K:F]\,|\,nc$, where $c:=\mathrm{c}(\Delta\otimes K)$. If K splits Δ, then $c=\deg(\Delta)$ and hence $[K:F]=n\deg(\Delta)=nc$. Therefore, $\mathrm{Hom}_F(K,A)$ is non-empty. Suppose $[K:F]|nc$. Then $\deg(\Delta)|c$ and $\deg(\Delta)=c$. This shows that K splits Δ.

(2) Suppose there is an embedding of K in A. Then there are positive integers n_i with $n=\sum n_i$ and there is an embedding of K_i in $\mathrm{Mat}_{n_i}(\Delta)$. Since $[K_i:\ F]\,|\,n_i\deg(\Delta)$, it follows from

$$[K:F]=\deg(A)=\sum_i n_i\deg(\Delta)\geq\sum_i [K_i:F]=[K:F]$$

that $[K_i:F]=n_i\deg(\Delta)$ for each $i=1,\dots,s$. Therefore, K_i splits Δ. Conversely, if K_i splits Δ for each i, then $[K_i:F]=m_i\deg(\Delta)$ for a positive integer m_i and $c_i:=\mathrm{c}(\Delta\otimes_F K_i)=\deg(\Delta)$. Then we have

$$n=\sum_i m_i, \quad \text{and} \quad [K_i:F]\,|\,m_i c_i, \quad \forall\, i=1,\dots,s.$$

It follows from Lemma 2.4 that there is an embedding of K into A. ∎

3. Numerical solutions to (Q1) and (Q2)

Let the ground field F be a local field in § 3.1 and be a global field from § 3.2 on. We let A denote a central simple algebra over F and K a commutative semi-simple algebra over F, and study the embedding of K into A over F.

3.1. **Local results.** Let F denote a local field.

Lemma 3.1. *Let $A=\mathrm{End}_\Delta(V)=\mathrm{Mat}_n(\Delta)$ be a central simple algebra over F.*

(1) *Let K be a finite field extension of F. The following statement are equivalent:*
    (a) *There exists an embedding of K in A over F.*
    (b) $[K:F]\,|\,n\cdot\mathrm{c}(\Delta\otimes_F K)$.
    (c) $[K:F]\,|\,n\deg(\Delta)$.

(2) *Let $K=\prod_{i=1}^{s}K_i$ be a commutative semi-simple algebra over F. Then there exists an embedding of K in A if and only if there are positive integers n_i for $i=1,\dots,s$ such that*

$$n=\sum_{i=1}^{s} n_i, \quad \textit{and} \quad [K_i:F]\,|\,n_i\deg(\Delta), \quad \forall\, i=1,\dots,s. \tag{3.1}$$

It follows from Lemma 2.4 that the statements (a) and (b) are equivalent. The implication $(b) \implies (c)$ is trivial. Put $\delta := \deg(\Delta)$ and $k := [K : F]$. If $\mathrm{inv}(\Delta) = a/\delta$ with $(a, \delta) = 1$, then (see [8])

$$\mathrm{inv}(\Delta \otimes_F K) = [K : F]\, \mathrm{inv}(\Delta) = \frac{ak}{\delta} = \frac{a'}{\delta'}, \quad \text{with} \ \ (a', \delta') = 1,$$

where $\delta = \delta' c$, $ak = a'c$, and $c := (k, \delta)$. It follows that

$$\mathrm{c}(\Delta \otimes_F K) = ([K : F], \deg(\Delta)). \tag{3.2}$$

Note that $(\delta', k) = 1$, so we have

$$k \,|\, n\delta \iff k \,|\, nc\delta' \iff k \,|\, nc.$$

The statement Lemma 3.1 (2) follows from Lemma 2.4 and Lemma 3.1 (1).

Now consider the case where $K = \prod_{i=1}^{s} K_i$ is a commutative semi-simple F-algebra. Put

$$c_i := ([K_i : F], \deg(\Delta)), \quad \ell_i := [K_i : F]/c_i. \tag{3.3}$$

For any positive integer n_i, we have

$$[K_i : F] \,|\, n_i \deg(\Delta) \iff \ell_i \,|\, n_i. \tag{3.4}$$

Put

$$\mathcal{E}_F(K, A) := \{x = (x_1, \dots, x_s) \in \mathbb{N}^s \mid \sum_{i=1}^{s} \ell_i x_i = \dim_\Delta V\}. \tag{3.5}$$

If a tuple $\mathbf{n} = (n_1, \dots, n_s)$ is a solution to (3.1), then the tuple $x = (x_1, \dots, x_s)$, where $x_i := n_i/\ell_i$, is an element in $\mathcal{E}_F(K, A)$. Conversely, any element x in $\mathcal{E}_F(K, A)$ gives a solution $\mathbf{n}$ to (3.1) by setting $n_i = \ell_i x_i$. Recall that $A^\times \backslash \mathrm{Hom}_F^*(K, A)$ is the set of equivalence classes of embeddings of F-algebras from K into A.

Proposition 3.2. *There is a natural bijection*

$$e : A^\times \backslash \mathrm{Hom}_F^*(K, A) \xrightarrow{\sim} \mathcal{E}_F(K, A). \tag{3.6}$$

Proof. Let φ and φ' be two maps in $\mathrm{Hom}_F^*(K, A)$, and let V_φ and $V_{\varphi'}$ be the induced (K, Δ)-bimodule structures on V. Write

$$V_\varphi = V_1 \oplus \cdots \oplus V_s \quad \text{and} \quad V_{\varphi'} = V_1' \oplus \cdots \oplus V_s',$$

where V_i and V_i' are (K_i, Δ)-bimodules. We have shown (Lemma 2.5) that φ and φ' are equivalent if and only if V_φ and $V_{\varphi'}$ are isomorphic as (K, Δ)-bimodules, equivalently, $V_i \simeq V_i'$ as (K_i, Δ)-bimodules for $i = 1, \dots, s$. Since each $\Delta \otimes_F K_i$ is simple, the latter is the same as the condition $\dim_\Delta V_i = \dim_\Delta V_i'$ for $i = 1, \dots, s$. One associates to φ a tuple

$$\mathbf{n} = (\dim_\Delta V_1, \dots, \dim_\Delta V_s)$$

which satisfies the condition (3.1) and determines the map φ up to equivalence. As such tuples are one-to-one in correspondence with elements in $\mathcal{E}_F(K, A)$. Then we show a bijection map $e : A^\times \backslash \mathrm{Hom}_F^*(K, A) \to \mathcal{E}_F(K, A)$ which is given by

$$e(\varphi) = (\dim_\Delta V_1/\ell_1, \dots, \dim_\Delta V_s/\ell_s). \tag{3.7}$$

This completes the proof of the proposition. ∎

3.2. **Global results.** From now on, we let F be a global field. Let A be a central simple algebra over F and K a finite field extension over F. We use the following notations.

- $A = \mathrm{End}_\Delta(V)$, where Δ is the division part of A, and V is a finite right Δ-module of rank n.
- $k := [K : F]$ and $\delta_0 := \deg(\Delta)$.
- For any place v of F, denote by F_v the completion of F at v. Put
$$K_v := K \otimes F_v = \prod_{w|v} K_w, \quad A_v := A \otimes F_v, \quad \Delta_v = \Delta \otimes F_v = \mathrm{Mat}_{s_v}(D_v),$$
where D_v is the division part of the central simple algebra Δ_v (we do not use the letter D as an algebra over F in this section; do not confuse D_v as the completion of D) and s_v is the capacity of Δ_v.
- $k_w := [K_w : F_v]$ and $d_v := \deg(D_v)$, where w is a place of K over v.
- $\Delta \otimes_F K = \mathrm{Mat}_c(\Delta')$ and $\delta' := \deg(\Delta')$, where Δ' is the division part of the central simple algebra $\Delta \otimes K$ over K, and c is its capacity. One has
$$\delta_0 = \delta' c. \tag{3.8}$$
- For any place w of K, put
$$\Delta'_w := \Delta' \otimes_K K_w = \mathrm{Mat}_{t_w}(D'_w), \quad d'_w := \deg(D'_w),$$
where D'_w is the division part of the central simple algebra Δ'_w and t_w is the local capacity of Δ' at w.
- $c_w := \mathrm{c}(D_v \otimes_{F_v} K_w)$, i.e. $D_v \otimes_{F_v} K_w = \mathrm{Mat}_{c_w}(D'_w)$. One has
$$d_v = d'_w c_w. \tag{3.9}$$
It follows from
$$\Delta \otimes_F K_w = (\Delta \otimes F_v) \otimes_{F_v} K_w = \Delta_v \otimes K_w = \mathrm{Mat}_{s_v c_w}(D'_w) \quad \text{and}$$
$$\Delta \otimes_F K_w = (\Delta \otimes_F K) \otimes_K K_w = \mathrm{Mat}_c(\Delta') \otimes_K K_w = \mathrm{Mat}_{ct_w}(D'_w)$$
that
$$s_v\, c_w = c\, t_w. \tag{3.10}$$
- For any rational number $a \in \mathbb{Q}$, we write $\mathbf{d}(a)$ for the positive denominator of a in its reduced form, and $\mathbf{n}(a)$ for its numerator.
- For each place v of F, write
$$\mathrm{inv}_v(\Delta) = \frac{a_v}{\delta_0} = \frac{a'_v\, s_v}{d_v\, s_v} = \frac{a'_v}{d_v}, \quad (a'_v, d_v) = 1 \text{ and } s_v = (a_v, \delta_0).$$
One has, by the Grunwald-Wang theorem
$$\delta_0 = \mathrm{lcm}\{d_v\}_{v \in V^F} \quad \text{and} \quad (\gcd\{a_v\}_{v \in V^F}, \delta_0) = 1, \tag{3.11}$$
where V^F denotes the set of all places of F.
- For each place w of K, write
$$\mathrm{inv}_w(\Delta') = \frac{b_w}{\delta'} = \frac{b'{}_w\, t_w}{d'_w\, t_w} = \frac{b'{}_w}{d'_w}, \quad (b'{}_w, d'_w) = 1 \text{ and } t_w = (b_w, \delta').$$
One has
$$\delta' = \mathrm{lcm}\{d'_w\}_{w \in V^K} \quad \text{and} \quad (\gcd\{b_w\}_{w \in V^K}, \delta') = 1, \tag{3.12}$$
where V^K denotes the set of all places of K.

- It follows from $\mathrm{inv}(D'_w)=\mathrm{inv}(D_v)[K_w:F_v]$ (see [8]) that

$$c_w=(d_v,k_w). \tag{3.13}$$

Given K and A, we have, for each place v of F,

- a tuple $(k_w)_{w|v}$ of positive integers, and
- a rational number $\mathrm{inv}_v(\Delta)=a'_v/d_v$

satisfying the following conditions:

(a) $\sum_{w|v}k_w=k$ for all $v\in V^F$,

(b) (i) $d_v=1$ if v is a complex place,
  (ii) $d_v\in\{1,2\}$ if v is a real place,
  (iii) $d_v=1$ for almost all v, and
  (iv) (Global class field theory) one has

$$\sum_{v\in V^F}\frac{a'{}_v}{d_v}=0 \quad (\text{ in } \mathbb{Q}/\mathbb{Z}).$$

We compute all other numerical invariants δ_0, c_w, d'_w, δ' and c as follows.

(i) The (global) degree δ_0 of Δ can be computed by (3.11).
(ii) Then one computes the local capacity c_w of $D_v\otimes_{F_v}K_w$ and the (local) degree d'_w of D'_w by (3.13) and (3.9), respectively.
(iii) Using (3.12) we compute the (global) degree δ' of Δ' and then compute the (global) capacity c of $\Delta\otimes K$ using (3.8).

We define the following condition (G stands for “global”)

(**G**) $k\,|\,n\,c$.

Proposition 3.3. *The set* $\mathrm{Hom}_F(K,A)$ *is non-empty if and only if the condition* (**G**) *holds.*

Proof. This follows from Lemma 2.4. ∎

Now we formulate the corresponding local conditions. Note that

$$K_v=\prod_{w|v}K_w \quad\text{and}\quad A_v=\mathrm{Mat}_{ns_v}(D_v).$$

Put

$$\mathcal{E}:=A^\times\backslash\,\mathrm{Hom}_F(K,A); \tag{3.14}$$

the Noether-Skolem theorem says that if this set is non-empty then it has one element. For each place v of F, define a set (c.f. (3.5))

$$\mathcal{E}_v:=\mathcal{E}_{F_v}(K_v,A_v)=\{(x_w)_{w|v}\,|\,x_w\in\mathbb{N},\ \sum_{w|v}\ell_w x_w=ns_v\}, \tag{3.15}$$

where $\ell_w:=k_w/c_w$. Define the following condition (L stands for “local”)

(**L**) The set $\mathcal{E}_v$ is non-empty for all $v\in V^F$.

Proposition 3.4. *There is an embedding of K_v in A_v over F_v if and only if the set $\mathcal{E}_v$ is non-empty.*

Proof. This follows from Proposition 3.2. ∎

We have the following implication

$$\text{the condition } (\mathbf{G}) \text{ holds} \implies \text{the condition } (\mathbf{L}) \text{ holds}.$$

The local-global principle then asks whether the converse is also true.

3.3. Special vectors and the local-global principle.

Let

$$e_v : A_v^\times \backslash \mathrm{Hom}^*_{F_v}(K_v, A_v) \xrightarrow{\sim} \mathcal{E}_v$$

be the corresponding bijection obtained in Proposition 3.2. Let us suppose first that the set $\mathrm{Hom}_F(K,A) = \mathrm{Hom}^*_F(K,A)$ of embeddings of K in A over F is *non-empty.* For any element φ in $\mathrm{Hom}_F(K,A)$, let $\varphi_v \in \mathrm{Hom}_{F_v}(K_v, A_v)$ be the extension of φ by F_v-linearity, and let $[\varphi_v]$ be its equivalence class. Then one defines an element $\mathbf{x}_v \in \mathcal{E}_v$ by

$$\mathbf{x}_v := e_v([\varphi_v]).$$

The association $\varphi \mapsto \mathbf{x}_v$ induces a well-defined map, which we denote again by e_v,

$$e_v : \mathcal{E} \to \mathcal{E}_v.$$

The non-emptiness of $\mathrm{Hom}_F(K,A)$ implies the existence of such a vector $\mathbf{x}_v$ in $\mathcal{E}_v$ for each place $v \in V^F$. We now calculate these special vectors explicitly.

The map φ gives rise to a (K, Δ)-bimodule structure on V. Since V is free K-module of rank $n\delta_0^2/k$, its completion $V \otimes_F F_v$ is also a free K_v-module of same rank. Therefore, one has the decomposition

$$V \otimes F_v = \bigoplus_{w|v} V_w,$$

where each factor V_w is a $(K_w, \mathrm{Mat}_{s_v}(D_v))$-bimodule of K_w-rank $n\delta_0^2/k$ (recall that $\Delta_v = \mathrm{Mat}_{s_v}(D_v)$). Using the Morita equivalence, the module V_w is isomorphic to $W_w^{\oplus s_v}$ for a (K_w, D_v)-bimodule W_w of D_v-rank $ns_v k_w/k$. Using the formula (3.7), the w-component $\mathbf{x}_w$ of the vector $\mathbf{x}_v$ is given by

$$\mathbf{x}_w := \dim_{D_v} W_w/\ell_w = ns_v c_w/k, \tag{3.16}$$

which is a positive integer. Recall that $c_w = (k_w, d_v)$ and $\ell_w = k_w/c_w$.

Therefore, this leads us to the following definition of *special vectors* no matter the set $\mathcal{E}$ is non-empty or not. For each place v of F we define a vector (still denoted by) $\mathbf{x}_v = (\mathbf{x}_w)_{w|v} \in \prod_{w|v} \mathbb{Q}_{>0}$ by (3.16), and we call them *special vectors.* The above calculation shows if the set $\mathcal{E}$ is non-empty, then the vector $\mathbf{x}_v$ is the image of the map e_v.

Proposition 3.5. *Notations as above. If the set $\mathcal{E}$ is non-empty, then one has*

$$\mathbf{x}_v \in \mathcal{E}_v, \quad \forall\, v \in V^F,$$

or equivalently, each vector $\mathbf{x}_v$ lies in $\prod_{w|v} \mathbb{N}$ for all $v \in V^F$.

If we denote by $\bar{\mathbf{x}}_w$ the class of $\mathbf{x}_w$ in $\mathbb{Q}/\mathbb{Z}$, then we associate to the pair (K,A) an element

$$\bar{\mathbf{x}} = (\bar{\mathbf{x}}_w)_{w \in V^K} \in \bigoplus_{w \in V^K} \mathbb{Q}/\mathbb{Z}. \tag{3.17}$$

Then Proposition 3.5 states that the vanishing of the class $\bar{\mathbf{x}}$ is a necessary condition for the set $\mathcal{E}$ to be non-empty. The following result states that this is the only obstruction to the local-global principle.

Theorem 3.6. *Notations as above. We have*

$$\mathrm{Hom}_F^*(K, A) \neq \emptyset \iff \bar{\mathbf{x}} = 0.$$

Proof. Note that the condition $\bar{\mathbf{x}} = 0$ implies $\mathbf{x}_v \in \mathcal{E}_v$ and hence $\mathrm{Hom}_{F_v}^*(K_v, A_v)$ is non-empty for all $v \in V^F$. The implication $\Longrightarrow$ is already proved. To show the other direction, we must show that the condition $(\mathbf{G})$ $k \mid nc$ holds. Using $c = \delta_0/\delta'$ and $\delta' = \mathrm{lcm}\{d'_w\}$, we rewrite the condition $(\mathbf{G})$ as

$$k \mid (n\delta_0/d'_w), \quad \forall\, w \in V^K. \tag{3.18}$$

Using (3.9) and (3.16), we have

$$\mathbf{x}_w = ns_v c_w/k = ns_v d_v/kd'_w = n\delta_0/kd'_w \in \mathbb{N}$$

for all $w \in V^K$. This verifies the condition $(\mathbf{G})$ and hence proves the theorem. ∎

3.4. **Examples.** We will show an example of a pair (K, A), where K/F is a Galois extension and A is a central simple F-algebra so that

- the set $\mathrm{Hom}_{F_v}^*(K_v, A_v)$ is non-empty for all $v \in V^F$, and
- the set $\mathrm{Hom}_F(K, A)$ is empty.

This particularly shows that the question (Q3) has a negative answer. Let K/F is a Galois extension of degree 8. Choose two places v_1 and v_2 of F so that $k_w = 2$ for all $w|v_1$ or $w|v_2$. Such places exist by the Chebotarev density theorem. Let A be a central simple algebra over F of degree 24 that is ramified exactly at the two places v_1 and v_2, and

$$A_{v_1} = \mathrm{Mat}_6(D_{v_1}), \quad \text{and} \quad A_{v_2} = \mathrm{Mat}_6(D_{v_2}),$$

where D_{v_1} and D_{v_2} are central division algebras of degree 4. The existence of such an A follows from the global class field theory. One has $c_w = 2$ and $\ell_w = 1$. The sets $\mathcal{E}_v$ for unramified places v are non-empty. For $v = v_i$, $i = 1, 2$, One see that

$$\mathcal{E}_{v_i} = \{(x_i) \in \mathbb{N}^4 | \sum_{i=1}^4 x_i = 6\} \neq \emptyset.$$

By Proposition 3.4, the sets $\mathrm{Hom}_{F_v}^*(K_v, A_v)$ are non-empty for all $v \in V^F$. On the other hand, $\mathbf{x}_w = 6c_w/k = 12/8$ for any $w|v_1$ or $w|v_2$, which is not an integer. By Theorem 3.6, the set $\mathrm{Hom}_F(K, A)$ is empty.

The same argument shows that there is a Galois extension K/F of degree p^m, and a central simple algebra A/F of degree $p^m q$, where p and q are primes, with $m \geq 2$ and $p < q$, so that the local-global principle for embedding K in A fails. One takes $k_w = p$ and $d_v = p^2$. Then $c_w = p$ and $\ell_w = 1$, We see that $\mathcal{E}_{v_i}$ is non-empty from the inequality $p^{m-1} \leq p^{m-2}q$. However, $\mathbf{x}_w = p^{m-2}qp/p^m \notin \mathbb{N}$.

4. The local-global principle

Keep the notation in § 3.2 and 3.3. In this section, we study on the local-global principle in detail.

For the convenience of discussion, we make the following definitions.

Definition 4.1.

(1) Let K be a finite field extension over a global field F and A be a central simple algebra over F. We say that the Hasse principle for the pair (K, A) holds if the following conditions are equivalent:

$$\mathrm{Hom}^*_{F_v}(K_v, A_v) \neq \emptyset, \quad \forall\, v \in V^F \Longleftrightarrow \mathrm{Hom}^*_F(K, A) \neq \emptyset.$$

In other words, the conditions $(\mathbf{G})$ and $(\mathbf{L})$ in § 3.2 are equivalent.

(2) Let (k, δ) be a pair of two positive integers with $k \mid \delta$. We say that the Hasse principle for the pair (k, δ) holds if for any finite field extension K over F of degree k and for any central simple algebra A over F of degree $\deg(A) = \delta$, the Hasse principle for the pair (K, A) holds.

Theorem 3.6 gives us an effective way to check the Hasse principle. Namely, we need to check

$$\mathbf{x}_w = ns_v c_w/k \overset{?}{\in} \mathbb{N}, \quad \forall\, w \in V^K \tag{4.1}$$

for a given pair (K, A).

4.1. Basic positive results.

Proposition 4.2. *Let A be a central simple algebra over F with index $\mathrm{i}(A) = \delta_0$ and degree $\deg A = n\delta_0$, and let K be a finite field extension over F of degree $[K : F] \mid \deg A$. Suppose one of the following properties holds:*

(1) *$[K : F] = \deg A$.*
(2) *K splits A.*
(3) *For any place v, the algebra A_v is either a division algebra or a matrix algebra over F_v.*

Then the Hasse principle for the pair (K, A) holds.

Proof.

(1) This is a well-known result; see Pierce [4, §18.4] and Prasad-Rapinchuk [5, Proposition A.1].

(2) Since K splits A, we have $d'_w = 1$ for all $w \in V^K$ and hence $\delta' = \underset{w}{\mathrm{lcm}}\{d'_w\} = 1$ and $c = \delta_0$. Thus,

$$k \mid n\delta_0 \Longleftrightarrow k \mid nc.$$

This verifies the condition $(\mathbf{G})$ and hence $\mathrm{Hom}_F(K, A) \neq \emptyset$.

(3) Suppose there exists an embedding of K_v in A_v for all $v \in V^F$. If A_v is a central division algebra, then $K_v = K_w$ is a field extension over F_v with $[K_w : F_v] = k$ and $c_w = k$. Then we have

$$\mathbf{x}_w = ns_v c_w/k = n \in \mathbb{N}, \quad \forall\, w \mid v.$$

For the other case that A_v is a matrix algebra over F_v, we have $s_v = \delta_0$. Thus, by the assumption $k \mid n\delta_0$, we have

$$\mathbf{x}_w = ns_v c_w/k = n\delta_0 c_w/k \in \mathbb{N}, \quad \forall\, w \mid v.$$

By Theorem 3.6, the set $\mathrm{Hom}_F(K,A)$ is not empty. ∎

4.2. **Construction of counterexamples.** Let K be a finite field extension over F with $[K:F]=k=p_1^{m_1}\cdots p_r^{m_r}$, and A be a central division algebra over F of degree $\delta=p_1^{n_1}\cdots p_r^{n_r}$, where p_i is a prime number, $n_i\in\mathbb{N}$, and $m_i\in\mathbb{Z}_{\geq 0}$ with $m_i\leq n_i$ for $i=1,\ldots,r$.

We shall construct examples for which the Hasse principle does not hold. Recall (Definition 4.1) that the Hasse principle for a pair (k,δ) does not hold if there exists a pair (K,A) such that $\mathrm{Hom}^*_{F_v}(K_v,A_v)\neq\emptyset$ for all $v\in V^F$ but $\mathrm{Hom}_F(K,A)=\emptyset$.

The construction is to set local data s_v, d_v, and a partition $k=\sum_{w|v}k_w$ of the integer k for some place v of F so that some w-component $\mathbf{x}_w$ of the special vector $\mathbf{x}_v$ is not integral. Then we apply Theorem 3.6 to conclude that the Hasse principle for certain pair (K,A) does not hold. To ensure that the data $k=\sum_{w|v}k_w$ come from a global field K, we need the following result.

Lemma 4.3 (c.f. [10, Lemma 3.2])**.** *Let S be a finite subset of V^F. Let L_v, for each $v\in S$, be any etale algebra over F_v of same degree $[L_v:F_v]=n$. Then there is a finite separable field extension K over F of degree n such that $K\otimes_F F_v\simeq L_v$ for all $v\in S$.*

Moreover, one also needs to ensure that the data s_v and d_v come from a central division algebra A over F. For this, we need the global class field theory; see [7, § 32] or [1].

Theorem 4.4. *Let S be a finite subset of V^F. For any positive integer δ, suppose we are given any set of rational numbers $\{a_v/d_v\}_{v\in S}$ with $(a_v,d_v)=1$ such that*

1. $\underset{v\in S}{\mathrm{lcm}}\{d_v\}=\delta$.
2. $\sum\limits_{v\in S}a_v/d_v=0\in\mathbb{Q}/\mathbb{Z}$.
3. $d_v=1$ *if v is complex.*
4. $d_v=1$ *or* 2 *if v is real.*

Then up to isomorphism, there is a unique central division algebra A over F with $\deg A=\delta$ such that $\mathrm{inv}\,A_v=a_v/d_v\in\mathbb{Q}/\mathbb{Z}$ for $v\in S$ and $\mathrm{inv}\,A_v=0\in\mathbb{Q}/\mathbb{Z}$ for $v\notin S$.

Let (k,δ) be a pair of positive integers with $k\mid\delta$. For any partition λ of k

$$\lambda=(k_1,\ldots,k_t),\quad k_1\leq\cdots\leq k_t,\quad \sum_{i=1}^t k_i=k$$

and a decomposition $\mu=(s,d)$ of $\delta=sd$ into the product of two positive integers, we set

$$\mathcal{E}(\lambda,\mu):=\{(x_i)\in\mathbb{N}^t\mid \sum_{i=1}^t \ell_i x_i=s\},$$

where $\ell_i:=k_i/(k_i,d)$, and define a vector $\mathbf{x}(\lambda,\mu)\in\mathbb{Q}^t$

$$\mathbf{x}(\lambda,\mu):=(\mathbf{x}_i),\quad \mathbf{x}_i=\mathbf{x}_i(\lambda,\mu):=s(k_i,d)/k,\quad \forall\, i=1\ldots,t.$$

Let $\mathrm{LD}(k,\delta)$ be the set of pairs (λ,μ) of partitions λ of k and a decomposition μ of δ such that $\mathcal{E}(\lambda,\mu)$ is non-empty ("LD" stands for all possible local decompositions).

We transform the problem of checking the Hasse principle for the pair (k,δ) in purely combinatorial terms.

Theorem 4.5. *Let (k,δ) be a pair of positive integers with $k \mid \delta$.*

1. *Suppose for all elements $(\lambda,\mu) \in \mathrm{LD}(k,\delta)$, the vector $\mathbf{x}(\lambda,\mu)$ has integral components. Then the Hasse principle for (k,δ) holds.*
2. *Suppose there is an element $(\lambda,\mu) \in \mathrm{LD}(k,\delta)$ such that $\mathbf{x}_i(\lambda,\mu) \notin \mathbb{N}$ for some i. Then there are a finite separable field extension K of degree k over F and a central division algebra A of degree δ over F so that the Hasse principle for (K,A) fails. In particular, the Hasse principle for the pair (k,δ) fails.*

Proof. (1) Let K be any finite field extension of F of degree k and A be any central simple algebra over F of degree δ. For any place v of F, we get a partition $\lambda = (k_w)_{w|v}$ of k and a decomposition $\mu = (s,d)$ of δ, where s is the capacity of $A_v = A \otimes F_v$ and d is the index of A_v. The assumption $\mathrm{Hom}^*_{F_v}(K_v, A_v) \neq \emptyset$ assures that $\mathcal{E}(\lambda,\mu) = \mathcal{E}_v$ is non-empty. Therefore, the pair (λ,μ) is an element in $\mathrm{LD}(k,\delta)$. Then the assumption gives that $\mathbf{x}_v = \mathbf{x}(\lambda,\mu)$ has integral components. This works for all $v \in V^F$. By Theorem 3.6, we have $\mathrm{Hom}^*_F(K,A) \neq \emptyset$.

(2) Let $(\lambda,\mu) \in \mathrm{LD}(k,\delta)$, where $\lambda = (k_1,\dots,k_t)$ and $\mu = (s,d)$, be an element such that $\mathbf{x}(\lambda,\mu)$ is not an integral vector. Consider a finite set $S = \{v_1, v_1', v_2, v_2'\}$ of 4 finite places of F. Let A be the central division algebra over F of degree δ with following local invariants (Theorem 4.4):

- $\mathrm{inv}_{v_1}(A) = -\mathrm{inv}_{v_1'}(A) = 1/d$ and $\mathrm{inv}_{v_2}(A) = -\mathrm{inv}_{v_2'}(A) = 1/\delta$.
- $\mathrm{inv}_v(A) = 0$ for all $v \notin S$.

Let L_{v_2} and $L_{v_2'}$ be any finite separable *field* extensions of degree k over F_{v_2} and $F_{v_2'}$, respectively. Let $L_{v_1} = \prod_{i=1}^t E_i$ and $L_{v_1'} = \prod_{i=1}^t E_i'$ where E_i (resp. E_i') is a separable field extension of F_{v_1} (resp. of $F_{v_1'}$) of degree k_i for all i. By Lemma 4.3, there exists a finite separable field extension K of F of degree k such that $K \otimes F_v \simeq L_v$ for all $v \in S$. It follows from $(\lambda,\mu) \in \mathrm{LD}(k,\delta)$ that $\mathcal{E}_{v_1}$ and $\mathcal{E}_{v_1'}$ are non-empty. Also the sets $\mathcal{E}_{v_2}$ and $\mathcal{E}_{v_2'}$ are non-empty as K_{v_2} and $K_{v_2'}$ are fields. Since A_v is a matrix algebra for $v \notin S$, these sets $\mathcal{E}_v$ are non-empty, too. Finally, since $\mathbf{x}(\lambda,\mu)$ is not integral, the special vector $\mathbf{x}_{v_1} = \mathbf{x}(\lambda,\mu)$ is not integral. Therefore, we have constructed a pair (K,A) for which the Hasse principle fails. ∎

Proposition 4.6. *Let $\delta = p_1^{n_1} \cdots p_r^{n_r}$, and let k be a positive integer with $k > 1$ and $k \mid \delta$, where each p_i is a prime number. Assume that $k \le \delta/p_i^{n_i}$ and $p_i \mid k$ for some $1 \le i \le r$ (so $r \ge 2$). Then the Hasse principle for the pair (k,δ) does not hold.*

Proof. Without lose of generality, we may assume $i = 1$. We write $k = p_1^{m_1} k'$ for some prime-to-p_1 integer k'. Let

(i) $s = \delta/p_1^{n_1} = p_2^{n_2} \cdots p_r^{n_r}$.
(ii) $\lambda = (1,\dots,1) =: (1^{\oplus k})$.

Then one has

$$d = p_1^{n_1}, \quad c_i := (k_i, d) = 1, \quad \ell_i := k_i/c_i = 1,$$

and

$$\mathbf{x}_i = s/k = p_2^{n_2} \cdots p_r^{n_r}/p_1^{m_1} k' \notin \mathbb{N}, \quad \forall i.$$

Note that the condition $k \le \delta/p_1^{n_1} = s$ implies the equation $X_1 + \cdots + X_k = s$ has a positive integral solution, i.e. the set $\mathcal{E}(\lambda,\mu)$ is non-empty. By Theorem 4.5, the Hasse principle does not hold. ∎

Corollary 4.7. *Let δ be a positive integer divisible by two primes. Then there is a positive integer k with $k \mid \delta$ such that the Hasse principle for the pair (k, δ) does not hold.*

PROOF. Write $\delta = p_1^{n_1} \cdots p_r^{n_r}$, $r \geq 2$, and assume that p_1 is the smallest prime divisor. Let $k = p_1$. Then the corollary follows from Proposition 4.6. ∎

Corollary 4.8. *Let δ be a positive integer divisible by two primes. Then there are a central simple algebra A of degree δ and a finite field extension of F of degree $[K : F] \mid \delta$ so that the Hasse principle for the pair (K, A) does not hold.*

Proposition 4.6 tells us that when δ is divisible by two primes and k is "small" comparable to δ, the Hasse principle for (δ, k) fails. It is also well-known that when $k = \delta$ the Hasse principle holds. Therefore, it is natural to ask the Hasse principle when δ is a prime power or k is very close to δ. So we consider the following two more questions:

(Q4) Does the Hasse principle for a pair (k, δ) hold if $\delta = p^n$ is a prime power?
(Q5) Does the Hasse principle for a pair (k, δ) hold if δ / k is a prime?

5. ANSWERS FOR (Q4) AND (Q5)

5.1. **A reduction step.** Keep the notation in the previous section. Write $k = p_1^{m_1} \cdots p_r^{m_r}$ and $\delta = p_1^{n_1} \cdots p_r^{n_r}$, where p_i is a prime number and $0 \leq m_i \leq n_i \in \mathbb{N}$ for $i = 1, \dots, r$.

Lemma 5.1. *Let $\lambda = (k_1, \dots, k_t)$ be a partition of k and $\mu = (s, d)$ be a decomposition of δ. Put $h_i := k_i / \gcd(k_i, \delta)$ and*

$$\lambda' := ((k_i, \delta)^{\oplus h_i})_i = (k_{ij})_{1 \leq i \leq t, 1 \leq j \leq h_i},$$

where $k_{ij} := (k_i, \delta)$. Then

- *If $\mathcal{E}(\lambda, \mu) \neq \emptyset$, then $\mathcal{E}(\lambda', \mu) \neq \emptyset$.*
- *The vector $\mathbf{x}(\lambda, \mu)$ is not integral if and only if $\mathbf{x}(\lambda', \mu)$ is not integral.*

PROOF. (1) The condition $\mathcal{E}(\lambda, \mu) \neq \emptyset$ implies that the equation

$$\sum_{i=1}^{t} \ell_i X_i = s$$

has a positive integral solution, say (x_i), where $\ell_i = k_i/(k_i, d)$. We need to show that the equation

$$\sum_{i=1}^{t} \sum_{j=1}^{h_i} \ell_{ij} Y_{ij} = s$$

also has a positive integral solution, where $\ell_{ij} := k_{ij}/(k_{ij}, d) = (k_i, \delta)/\gcd(k_i, \delta, d) = (k_i, \delta)/(k_i, d)$. One has $\ell_i = \ell_{ij} h_i$ for all i. Put $Y_{ij} = x_i$, then the equation has a positive integral solution and hence the set $\mathcal{E}(\lambda', \mu)$ is non-empty.

(2) We have, for all i and j,

$$\mathbf{x}_i = s(k_i, d)/k = s \gcd(k_i, \delta, d)/k = s(k_{ij}, d)/k = \mathbf{x}_{ij}.$$

Therefore, the vector $\mathbf{x}(\lambda, \mu)$ is not integral if and only if $\mathbf{x}(\lambda', \mu)$ is not integral. This proves the lemma. ∎

Corollary 5.2. *Let (k,δ) be a pair as above. Suppose that for all $(\lambda,\mu)\in \mathrm{LD}(k,\delta)$ with $\lambda=(k_1,\dots,k_t)$ satisfying*

$$k_i \mid \delta, \quad \forall\, i=1,\dots,t \tag{5.1}$$

and that the vector $\mathbf{x}(\lambda,\mu)$ is integral. Then the Hasse principle for the pair (k,δ) holds.

Proof. This follows from Theorem 4.5 and Lemma 5.1. ∎

Corollary 5.2 tells us that if one wants to show that the Hasse principle for a pair (k,δ) fails, then it is enough to search pairs $(\lambda,\mu)\in\mathrm{LD}(k,\delta)$ with λ in the form (5.1) such that that the vector $\mathbf{x}(\lambda,\mu)$ is not integral. As we will see, it is actually easier to find a pair (λ,μ) with coordinates k_i of λ satisfying

$$k_i \mid k, \quad \forall\, i=1,\dots,t \tag{5.2}$$

with the desired property.

Put

$$I:=\{(a_1,\dots,a_r)\in\mathbb{Z}_{\ge 0}\mid 0\le a_i\le n_i \text{ for all } i\}.$$

Lemma 5.3. *There is a one-to-one correspondence between elements (λ,μ) in $\mathrm{LD}(k,\delta)$ with (5.1), and all non-negative integral solutions of the equations*

$$\begin{cases} \sum\limits_{(a_1,\dots,a_r)\in I} p_1^{a_1}\cdots p_r^{a_r} X_{(a_1,\dots,a_r)} = k \\ \sum\limits_{(a_1,\dots,a_r)\in I} \ell_{(a_1,\dots,a_r)}(Y_{(a_1,\cdots,a_r),1}+\cdots+Y_{(a_1,\dots,a_r),X_{(a_1,\dots,a_r)}}) = s \end{cases} \tag{5.3}$$

with the property that if $X_{(a_1,\dots,a_r)}>0$ then $Y_{(a_1,\cdots,a_r),j}>0$ for all $j=1,\dots,X_{(a_1,\dots,a_r)}$, where $(s,d)=\mu$ and $\ell_{(a_1,\dots,a_r)}=p_1^{a_1}\cdots p_r^{a_r}/(p_1^{a_1}\cdots p_r^{a_r},d)$.

Note that the number of variables of the second equation in (5.3) depends on the solutions of the first equation. This is not a standard system of equations, however, it is slightly easier for us to solve equations in this form. Solutions of the first equation correspond to partitions of k with the condition (5.1), and the second one is nothing but the defining equation for the set $\mathcal{E}(\lambda,\mu)$.

Now we use Corollary 5.2 and Lemma 5.3 to investigate the questions (Q4) and (Q5).

5.2. **Answers for (Q4).**

Theorem 5.4. *Let $(k,\delta)=(p^m,p^n)$, where p is a prime and $1\le m\le n\in\mathbb{N}$. The Hasse principle for the pair $(k,\delta)=(p^m,p^n)$ holds if and only if one of the following conditions holds:*

(i) *$m=n$.*
(ii) *$m=1$ or 2.*

The case (i) follows from Proposition 4.2 (1). Theorem 5.4 (ii) follows from Propositions 5.6 and 5.7 below.

Lemma 5.5. [4, § 13.4] *Let D be a central division algebra over any field F, and K/F be a field extension such that $[K:F]$ is a prime divisor of $\deg D$. The following properties are equivalent.*

(i) *K is isomorphic to a subfield of D.*
(ii) *$D\otimes K$ is not a division algebra.*

(iii) $\deg D = [K : F](\mathrm{i}(D \otimes K))$.

Proposition 5.6. *Let* (k, δ) *be as in Theorem 5.4. If* $m = 1$ *or* 2*, then the Hasse principle for the pair* (k, δ) *holds.*

Proof. (1) $m = 1$. Suppose there exists an embedding of K_v in A_v for all $v \in V^F$. By Lemma 5.5, it suffices to check that $A \otimes K$ is not a division algebra, i.e. $\delta' \neq p^n$. Since $\delta' = \mathrm{lcm}\{d'_w\}$, one needs to show $d'_w \neq p^n$ for all $w \in V^K$. If there exists $w \in V^K$ with $d'_w = p^n$, then $\deg A_v = p^n$ where $v \in V^F$ is the place with $w \mid v$. Suppose $K_v = K_{w_1} \times \cdots \times K_{w_r}$. By the assumption of the existence of local embeddings, we know that the equation $\ell_1 x_1 + \cdots + \ell_r x_r = 1$ has a positive integral solution, which implies that $r = 1$ and hence, $K_v = K_w$ is a field with $[K_w : F_v] = p$. Then $d'_w = p^n/[K_w : F_v] = p^{n-1}$, contradiction to $d'_w = p^n$. Thus $A \otimes K$ is not a division algebra.

(2) $m = 2$. In this case, it suffices to check $\mathbf{x}_i(\lambda, \mu) \in \mathbb{N}$ for all i and $(\lambda, \mu) \in \mathrm{LD}(p^2, p^n)$. If $s = p^a$ and $a \geq 2$, then $\mathbf{x}_i(\lambda, \mu) \in \mathbb{N}$ for all i. Therefore, one only needs to consider the following two cases:

(i) $s = 1$ and $d = p^n$. Since $(\lambda, \mu) \in \mathrm{LD}(p^2, p^n)$, the assumption $\mathcal{E}(\lambda, \mu) \neq \emptyset$ implies $\lambda = (k)$. Then $\mathbf{x}_1 = p^2/p^2 = 1 \in \mathbb{N}$. By Corollary 5.2, the Hasse principle holds.

(ii) $s = p$ and $d = p^{n-1}$. In the case of $n = 2$, this follows from Proposition 4.2 (1). So we may assume $n \geq 3$. By Lemma 5.3, one solves the system of linear equations

$$\begin{cases} X_0 + pX_1 + p^2 X_2 = p^2 \\ (Y_{0,1} + \cdots + Y_{0,X_0}) + (Y_{1,1} + \cdots + Y_{1,X_1}) + (Y_{2,1} + \cdots + Y_{2,X_2}) = p. \end{cases}$$

The solutions are $(X_0, X_1, X_2) = (0, 0, 1)$ and $(0, p, 0)$. Both solutions correspond to the partitions $\lambda_{(0,0,1)} = (p^2)$ and $\lambda_{(0,p,0)} = (p, \ldots, p) = p^{\oplus p}$ of k, respectively. Now one computes $\mathbf{x}_1(\lambda_{(0,0,1)}, \mu) = p \cdot p^2/p^2 = p$ and $\mathbf{x}_i(\lambda_{(0,1,0)}, \mu) = p \cdot p/p^2 = 1 \in \mathbb{N}$ for all i. By Corollary 5.2, the Hasse principle holds.

This proves the proposition. ∎

Proposition 5.7. *Let* (k, δ) *be as in Theorem 5.4. If* $m \geq 3$ *and* $m + 1 \leq n$*, then the Hasse principle for the pair* (k, δ) *fails.*

Proof. We consider a pair (λ, μ) as follows:

(i) $s = p^2$ and $d = p^{n-2}$.
(ii) $\lambda = (p^{m-3}, \ldots, p^{m-3}, p^{m-2}, \ldots, p^{m-2}, p^{m-1}, \ldots, p^{m-1})$
$= \big((p^{m-3})^{\oplus p},\ (p^{m-2})^{\oplus p-1},\ (p^{m-1})^{\oplus p-1}\big)$.

To check that the set $\mathcal{E}(\lambda, \mu)$ is non-empty, one solves the equation

$$(Y_{m-3,1} + \cdots + Y_{m-3,p}) + (Y_{m-2,1} + \cdots + Y_{m-2,p-1}) + (Y_{m-1,1} + \cdots + Y_{m-1,p-1}) = p^2.$$

Since the sum of coefficients $p + (p-1) + (p-1) = 3p - 2 \leq p^2$ for all primes p, this equation has at least one positive integral solution. Moreover, one computes $(p^{m-3}, d) = p^{m-3}$ and $\mathbf{x}_1(\lambda, \mu) = p^2 \cdot p^{m-3}/p^m = 1/p \notin \mathbb{N}$. By Theorem 4.5, the Hasse principle does not hold. This proves the proposition. ∎

5.3. **Answers for (Q5).** Let (k, δ) be a pair of positive integers such that $p_1 = \delta/k$ is a prime.

Proposition 5.8. *Suppose that p_1 is not the smallest prime divisor of δ. Then the Hasse principle for the pair (k, δ) does not hold.*

PROOF. Choose a prime divisor p_2 of δ with $p_2 < p_1$. We consider a pair (λ, μ) as follows:

(i) $s = p_1$ and $d = \delta/p_1 = k$.
(ii) $\lambda = (k/p_2, \dots, k/p_2) = (k/p_2)^{\oplus p_2}$.

To show that the set $\mathcal{E}(\lambda, \mu)$ is non-empty, one solves the equation

$$Y_{(n_1-1,n_2-1),1} + \cdots + Y_{(n_1-1,n_2-1),p_2} = p_1.$$

The condition $p_2 < p_1$ implies that the equation has at least one positive integral solution. In addition, one computes $(k/p_2, d) = k/p_2$ and $\mathbf{x}_i(\lambda, \mu) = (sk/p_2)/k = p_1/p_2 \notin \mathbb{N}$ for all i. By Theorem 4.5, the Hasse principle does not hold. ∎

The remaining case where p_1 is the smallest prime divisor is quite complicated. We restrict ourselves to the case where δ is divisible by exact two primes, i.e. $\delta = p_1^{n_1} p_2^{n_2}$, where p_1 and p_2 are primes. In this case, we give the complete answer. From now on, let $(k, \delta) = (p_1^{n_1-1} p_2^{n_2}, p_1^{n_1} p_2^{n_2})$ and we assume that $p_1 < p_2$.

Proposition 5.9. *Let $\delta = p_1 p_2$ and $k = \delta/p_1 = p_2$ with $p_1 < p_2$. Then the Hasse principle for the pair (k, δ) holds.*

PROOF. If $s = 1$, p_2, or $p_1 p_2$, then $\mathbf{x}_i(\lambda, \mu) \in \mathbb{N}$ for all λ and i. Therefore, it remains to check the case $s = p_1$ and $d = p_2$. In this case, the corresponding system of equations is

$$\begin{cases} X_{(0,0)} + p_1 X_{(1,0)} + p_2 X_{(0,1)} = p_2 \\ (Y_{(0,0),1} + \cdots + Y_{(0,0),X_{(0,0)}}) + p_1(Y_{(1,0),1} + \cdots + Y_{(1,0),X_{(1,0)}}) \\ \quad + (Y_{(0,1),1} + \cdots + Y_{(0,1),X_{(0,1)}}) = p_1. \end{cases}$$

Since $p_1 < p_2$, one sees that $(X_{(0,0)}, X_{(1,0)}, X_{(0,1)}) = (0, 0, 1)$ is the unique solution. It corresponds to the partition $\lambda = (p_2)$ of k. Then $(p_2, d) = p_2$ and $\mathbf{x}_1(\lambda, \mu) = p_1 \cdot p_2/p_2 = p_1 \in \mathbb{N}$. By Corollary 5.2, the Hasse principle for (k, δ) holds. This proves the proposition. ∎

Proposition 5.10. *Let $\delta = p_1^2 p_2$ and $k = \delta/p_1 = p_1 p_2$ with $p_1 < p_2$. Then the Hasse principle for the pair (k, δ) does not hold.*

PROOF. Since $p_1 < p_2$, we may choose two non-negative integers x_1, x_2 such that $p_2 - x_1 - p_1 x_2 = 1$ and $x_2 > 0$. Put $x_0 := p_1$. One checks that

$$x_0 + p_1 x_1 + p_1^2 x_2 = x_0 + p_1(p_2 - 1) = p_1 p_2 = k.$$

We consider a pair (λ, μ) as follows:

(i) $s = p_2$ and $d = p_1^2$.
(ii) $\lambda = \left(1^{\oplus x_0}, p_1^{\oplus x_1}, (p_1^2)^{\oplus x_2}\right)$.

Since components k_i of λ are $1, p_1$, or p_1^2, the coefficients ℓ_i of the equation for $\mathcal{E}(\lambda, \mu)$ are 1. To see $\mathcal{E}(\lambda, \mu)$ is non-empty, we check

$$x_0 + x_1 + x_2 = p_1 + (p_2 - 1 - p_1 x_2) + x_2 = p_2 + (p_1 - 1)(1 - x_2) \le p_2.$$

Now we check $\mathbf{x}_1(\lambda,\mu) = s(k_1,d)/k = p_2/p_1p_2 \notin \mathbb{N}$. By Theorem 4.5, the Hasse principle for the pair (k,δ) does not hold. ■

Proposition 5.11. *Let $\delta = p_1^{n_1}p_2^{n_2}$ and $k = \delta/p_1 = p_1^{n_1-1}p_2^{n_2}$ with $p_1 < p_2$ for n_1, $n_2 \geq 2$. Then the Hasse principle for the pair (k,δ) does not hold.*

Proof. We consider a pair (λ,μ) as follows:

(i) $s = \delta/p_2 = p_1^{n_1}p_2^{n_2-1}$ and $d = p_2$.

(ii) $\lambda = (p_1^{n_1-1},\dots,p_1^{n_1-1},p_2,\dots,p_2) = \left((p_1^{n_1-1})^{\oplus p_2^{n_2-1}}, p_2^{\oplus p_1^{n_1-1}p_2^{n_2-2}(p_2-1)}\right)$.

One looks at the equation

$$p_1^{n_1-1}(Y_{1,1}+\cdots+Y_{1,p_2^{n_2-1}}) + (Y_{2,1}+\cdots+Y_{2,p_1^{n_1-1}p_2^{n_2-2}(p_2-1)}) = p_1^{n_1}p_2^{n_2-1}.$$

This equation has a positive integral solution because we compare the sum of the coefficients with $p_1^{n_1}p_2^{n_2-1}$:

$$p_1^{n_1}p_2^{n_2-1} - p_1^{n_1-1}p_2^{n_2-1} - p_1^{n_1-1}p_2^{n_2-2}(p_2-1) = p_1^{n_1-1}p_2^{n_2-2}(p_1p_2 - 2p_2 + 1) \geq 0.$$

This shows $\mathcal{E}(\lambda,\mu) \neq \emptyset$. Moreover, one computes $(p_1^{n_1-1}, d) = 1$ and

$$\mathbf{x}_1(\lambda,\mu) = s/k = p_1^{n_1}p_2^{n_2-1}/p_1^{n_1-1}p_2^{n_2} = p_1/p_2 \notin \mathbb{N}.$$

By Theorem 4.5, the Hasse principle for (k,δ) does not hold. ■

Proposition 5.12. *Let $\delta = p_1^{n_1}p_2$ and $k = \delta/p_1 = p_1^{n_1-1}p_2$ with $p_1 < p_2$ for $n_1 \geq 3$. Then the Hasse principle for the pair (k,δ) does not hold.*

Proof. Case 1: $n_1 = 3$

One considers a pair (λ,μ) as follows:

(i) $s = p_1p_2$ and $d = p_1^2$.

(ii) $\lambda = (1,\dots,1,p_1,\dots,p_1,p_1^2) = \left(1^{\oplus p_1}, p_1^{\oplus p_1p_2-p_1-1}, p_1^2\right)$.

Similarly, one looks at the equation

$$(Y_{1,1}+\dots Y_{1,p_1}) + (Y_{2,1}+\cdots+Y_{2,p_1p_2-p_1-1}) + Y_3 = p_1p_2.$$

It follows from $p_1 + p_1p_2 - p_1 - 1 + 1 = p_1p_2$ that $\mathcal{E}(\lambda,\mu)$ is non-empty. Moreover, one computes $(1,d) = 1$ and $\mathbf{x}_1(\lambda,\mu) = s/k = p_1p_2/p_1^2p_2 = 1/p_1 \notin \mathbb{N}$.

Case 2: $n_1 \geq 4$

One considers a pair (λ,μ) as follows:

(i) $s = p_1^{n_1-2}$ and $d = p_1^2p_2$.

(ii) $\lambda = (p_2,\dots,p_2,p_1p_2,\dots,p_1p_2,p_1^2p_2,\dots,p_1^2p_2,\dots,p_1^{n_1-2}p_2,\dots,p_1^{n_1-2}p_2)$
$= \left(p_2^{\oplus p_1}, (p_1p_2)^{\oplus p_1-1}, (p_1^2p_2)^{\oplus p_1-1}, \dots, (p_1^{n_1-2}p_2)^{\oplus p_1-1}\right)$.

Similarly, by looking at the sum of coefficients of the defining equation for $\mathcal{E}(\lambda,\mu)$, we have

$$p_1 + (p_1-1) + \sum_{i=0}^{n_1-4} p_1^i(p_1-1) = p_1^{n_1-3} + 2p_1 - 2 \leq p_1^{n_1-2},$$

which implies that $\mathcal{E}(\lambda,\mu)$ is non-empty. Moreover, one computes $(p_2,d) = p_2$ and

$$\mathbf{x}_1(\lambda,\mu) = p_1^{n_1-2}p_2/k = p_1^{n_1-2}p_2/p_1^{n_1-1}p_2 = 1/p_1 \notin \mathbb{N}.$$

By Theorem 4.5, the Hasse principle for the pair (k,δ) does not hold. ■

Proposition 5.13. *Let* $\delta = p_1 p_2^{n_2}$ *and* $k = \delta/p_1 = p_2^{n_2}$ *with primes* $p_1 < p_2$ *and the integer* $n_2 \geq 2$. *Then the Hasse principle for the pair* (k, δ) *does not hold.*

Proof. One considers a pair (λ, μ) as follows:

(i) $s = p_1 p_2$ and $d = p_2^{n_2-1}$.
(ii) $\lambda = (p_2^{n_2-2}, \dots, p_2^{n_2-2}, p_2^{n_2-1}, \dots, p_2^{n_2-1}) = \big((p_2^{n_2-2})^{\oplus p_2}, (p_2^{n_2-1})^{\oplus p_2-1}\big)$.

Since $2p_2 - 1 \leq p_1 p_2$, one has $\mathcal{E}(\lambda, \mu) \neq \emptyset$. Moreover, one computes $(p_2^{n_2-1}, d) = p_2^{n_2-2}$ and

$$\mathbf{x}_1(\lambda, \mu) = p_1 p_2^{n_2-1}/k = p_1 p_2^{n_2-1}/p_2^{n_2} = p_1/p_2 \notin \mathbb{N}.$$

By Theorem 4.5, the Hasse principle for the pair (k, δ) does not hold. ∎

We summarize the results from Propositions 5.9 to 5.13 in the following theorem.

Theorem 5.14. *Let* $(\delta, k) = (p_1^{n_1} p_2^{n_2}, p_1^{n_1-1} p_2^{n_2})$, *where* $p_1 < p_2$ *are primes and* $n_1, n_2 \in \mathbb{N}$. *The Hasse principle for the pair* (k, δ) *holds if and only if* $(n_1, n_2) = (1, 1)$.

The following picture indicates for which pair (n_1, n_2) the Hasse principle for $(p_1^{n_1-1} p_2^{n_2}, p_1^{n_1} p_2^{n_2})$ with primes $p_1 < p_2$ holds.



We conclude this section by the following result.

Theorem 5.15. *The positive integers* δ *with the property that for any central simple algebra* A *over* F *of degree* δ *and any finite field extension* K *of degree* $[K : F]$ *dividing* δ *the Hasse principle for* (K, A) *holds are* $1, p, p^2, p^3$, *where* p *is a prime.*

Proof. This follows from Corollary 4.7 and Theorem 5.4. ∎

6. A GLUING RESULT OF EMBEDDINGS

In the section we study the embedding of a field extension in a simple algebra from the topological point of view. Let $A = \mathrm{Mat}_n(\Delta)$ be a central simple algebra over a global field F of degree m, where Δ is a central division algebra over F. Let K be a finite *separable* field extension over F of degree k dividing m.

We shall use freely the notations in the previous sections.

Let $\mathfrak{F}$ be a separable closed field which contains the fields F_v for all places $v \in V^F$. Recall that F_v denotes the completion of F at the place v. Note that such a big field $\mathfrak{F}$ exists. For example, let $\mathfrak{F}$ be a separable closure of the quotient Q of the big tensor $\otimes_{v \in V^F} F_v$ over F by a maximal ideal. Let F^{sep} and F_v^{sep} be the separable closures of F and F_v in $\mathfrak{F}$, respectively. Put $G_F := \mathrm{Gal}(F^{\mathrm{sep}}/F)$ and $G_{F_v} := \mathrm{Gal}(F_v^{\mathrm{sep}}/F_v)$; one has $G_{F_v} \subset G_F$. Let $G := \mathrm{Gal}(\mathfrak{F}/F)$ and $G_v := \mathrm{Gal}(\mathfrak{F}/F_v)$ for $v \in V^F$.

Definition 6.1. Let $\iota_v : K_v \hookrightarrow A_v$ be an F_v-embedding of K_v into A_v for each $v \in V^F$. Let

$$\iota_v \otimes \mathrm{id}_{\mathfrak{F}} : K_v \otimes_{F_v} \mathfrak{F} \hookrightarrow A_v \otimes_{F_v} \mathfrak{F}$$

be the induced $\mathfrak{F}$-embedding; note that $K_v \otimes_{F_v} \mathfrak{F} = K \otimes_F \mathfrak{F}$ and $A_v \otimes_{F_v} \mathfrak{F} = A \otimes_F \mathfrak{F}$. For any two places $v_1, v_2 \in V^F$, we say that ι_{v_1} and ι_{v_2} are $\mathfrak{F}$-equivalent, denoted by $\iota_{v_1} \sim_{\mathfrak{F}} \iota_{v_2}$ (or by $\iota_{v_1} \sim \iota_{v_2}$ for brevity if no confusion may cause) if the induced maps $\iota_{v_1} \otimes \mathrm{id}_{\mathfrak{F}}$ and $\iota_{v_2} \otimes \mathrm{id}_{\mathfrak{F}}$ are equivalent over $\mathfrak{F}$, that is, they are conjugate by an element in $(A \otimes_F \mathfrak{F})^\times$ (see Definition 2.2).

Suppose that we have an F-embedding $\iota : K \hookrightarrow A$. Let $\iota_v := \iota \otimes \mathrm{id}_{F_v}$ be the induced F_v-embedding. Then we have $\iota_{v_1} \otimes \mathrm{id}_{\mathfrak{F}} = \iota_{v_2} \otimes \mathrm{id}_{\mathfrak{F}}$, and in particular $\iota_{v_1} \sim \iota_{v_2}$ for any two places v_1 and v_2. Therefore, we may consider the following question*:

(Q6) Let $\iota_v : K_v \hookrightarrow A_v$ be an F_v-embedding of K_v into A_v for each $v \in V^F$. Suppose that $\iota_{v_1} \sim \iota_{v_2}$ for any two places $v_1, v_2 \in V^F$. Does there exist an F-embedding $\iota : K \hookrightarrow A$ such that $\iota \otimes \mathrm{id}_{F_v} \sim \iota_v$ (over F_v) for all $v \in V^F$?

The main result of this section is Theorem 6.7; which asserts that the question (Q6) has an affirmative answer.

Assume that $[K : F] = \deg A$. Suppose there is an F_v-embedding $\iota_v : K_v \hookrightarrow A_v$ for each $v \in V^F$. By a natural generalization of the Noether-Skolem theorem [6], any two $\mathfrak{F}$-embeddings $K \otimes_F \mathfrak{F} \hookrightarrow A \otimes_F \mathfrak{F}$ are conjugate to each other. By Theorem 6.7, then there is an F-embedding $\iota : K \hookrightarrow A$ so that $\iota \otimes \mathrm{id}_{F_v} \sim \iota_v$ for all $v \in V^F$. In particular, an F-embedding $K \hookrightarrow A$ exits. Therefore, Theorem 6.7 can be regarded as a generalization of the local-global principle for embeddings.

Recall (Definition 2.2) that

$$\mathcal{O}^*_{A_1,A_2} := A_2^\times \backslash \mathrm{Hom}^*_F(A_1, A_2)$$

denotes the set of equivalence classes of F-algebra embeddings of A_1 into A_2 for any two F-algebras A_1 and A_2. By Proposition 3.2, there is a natural bijection

$$\mathcal{O}^*_{K,A} \xrightarrow{\sim} \mathcal{E}_F(K, A). \tag{6.1}$$

Let

$$\Sigma := \mathrm{Hom}_F(K, \mathfrak{F}) = \{ \sigma_1, \dots, \sigma_k \} = \mathrm{Hom}_F(K, F^{\mathrm{sep}}).$$

We have $K \otimes_F \mathfrak{F} = \prod_{i=1}^k \mathfrak{F}_i$, where the field $\mathfrak{F}_i = \mathfrak{F}$ corresponds to the embedding σ_i for each i. As $\mathfrak{F}$ is separably closed, we have

$$\mathcal{O}^*_{K\otimes\mathfrak{F}, A\otimes\mathfrak{F}} = (\mathrm{Mat}_m(\mathfrak{F}))^\times \backslash \mathrm{Hom}^*_{\mathfrak{F}}(\prod_{i=1}^k \mathfrak{F}_i, \mathrm{Mat}_m(\mathfrak{F})).$$

By Proposition 3.2, we also get a bijection

$$\mathcal{O}^*_{K\otimes\mathfrak{F}, A\otimes\mathfrak{F}} \xrightarrow{\sim} \mathcal{E}_{\mathfrak{F}}(K \otimes \mathfrak{F}, A \otimes \mathfrak{F}) = \{ x = (x_1, \dots, x_k) \in \mathbb{N}^k \mid \sum_{i=1}^k x_i = m \}. \tag{6.2}$$

---

[*] This question was suggested by an referee. We thank him/her for the input

To each element $x=(x_1,\dots,x_k)\in\mathcal{E}_{\mathfrak{F}}(K\otimes\mathfrak{F},A\otimes\mathfrak{F})$, one associates a $\mathbb{N}$-valued function f on Σ by $f(\sigma_i):=x_i$. This gives a natural bijection between the set $\mathcal{E}_{\mathfrak{F}}(K\otimes\mathfrak{F},A\otimes\mathfrak{F})$ and the set

$$\mathcal{C}(K,A,\mathfrak{F}):=\{\,f:\mathrm{Hom}_F(K,\mathfrak{F})\longrightarrow\mathbb{N}\mid\ \sum_{i=1}^k f(\sigma_i)=m\,\}.$$

The set $\mathcal{C}(K,A,\mathfrak{F})$ has a natural G-action by $(\tau * f)(\sigma_i):=f(\tau\sigma_i)$ for $\tau\in G$ and $f\in\mathcal{C}(K,A,\mathfrak{F})$. The action of G on this set factors through the finite quotient $\mathrm{Gal}(E/F)$, where E is the Galois closure of K over F in F^{sep}, as we also have $\Sigma=\mathrm{Hom}_F(K,E)$.

Lemma 6.2.

1. *There exists a bijective map* $\phi:\mathcal{O}^*_{K\otimes\mathfrak{F},A\otimes\mathfrak{F}}\xrightarrow{\sim}\mathcal{C}(K,A,\mathfrak{F})$.
2. *The* G*-invariant subset* $\mathcal{C}(K,A,\mathfrak{F})^G$ *is equal to*
$$\{f:\Sigma\to\mathbb{N}\mid\ f(\sigma_i)=n/k,\forall\, i=1,\dots,k\}.$$
3. *There is a commutative diagram*
$$\begin{tikzcd}\mathcal{O}^*_{K,A} \arrow[r, "\otimes\,\mathrm{id}_{\mathfrak{F}}"] \arrow[d, "\psi"'] & \mathcal{O}^*_{K\otimes\mathfrak{F},A\otimes\mathfrak{F}} \arrow[d, "\phi"'] \\ \mathcal{C}(K,A,\mathfrak{F})^G \arrow[r] & \mathcal{C}(K,A,\mathfrak{F}),\end{tikzcd}$$
and the map ψ *is a bijection if the set* $\mathcal{O}^*_{K,A}$ *is non-empty.*

PROOF. (1) This follows from the natural bijection $\mathcal{E}_{\mathfrak{F}}(K\otimes\mathfrak{F},A\otimes\mathfrak{F})\simeq\mathcal{C}(K,A,\mathfrak{F})$ and the bijection (6.2).

(2) Since the action of G on Σ is transitive, the assertion follows.

(3) We may assume that the set $\mathcal{O}^*_{K,A}$ is non-empty. The proof is similar to that in Section 3.3. Let $[\varepsilon]\in\mathcal{O}^*_{K,A}$ be element. The map ε induces an embedding $\varepsilon\otimes\mathrm{id}_{\mathfrak{F}}:K\otimes_F\mathfrak{F}\hookrightarrow A\otimes_F\mathfrak{F}$, and by (1), it corresponds to a tuple $(x_1,\dots,x_k)$. Let $A=\mathrm{End}_\Delta(V)$ for a right Δ-vector space V of dimension n. Then V is a K-free $\Delta\otimes_F K$-module. As $(\Delta\otimes_F K)\otimes_F\mathfrak{F}=\prod_{i=1}^k\mathrm{Mat}_{\delta_0}(\mathfrak{F}_i)$, where $\delta_0=\deg(\Delta)$ and $\mathfrak{F}_i=\mathfrak{F}$, we have the decomposition $V\otimes_F\mathfrak{F}=\oplus_{i=1}^k V_i$, where V_i is a $\mathrm{Mat}_{\delta_0}(\mathfrak{F}_i)$-module. Note that $\dim_{\mathfrak{F}}V_i$ are the same for all i, and it is $m\delta_0/k$. Using the Morita equivalence, we have $V_i=W_i^{\oplus\delta_0}$ and

$$x_i:=\dim_{\mathfrak{F}}W_i=n\delta_0/k=m/k.$$

This completes the proof of the lemma. ∎

For each $v\in V^F$, let $K_v:=K\otimes_F F_v=\prod_{w|v}K_w$ and put $k_w:=[K_w:F_v]$. We have the following natural identifications

$$\Sigma=\mathrm{Hom}_F(K,F_v^{\mathrm{sep}})=\mathrm{Hom}_{F_v}(K_v,F_v^{\mathrm{sep}})=\coprod_{w|v}\Sigma_w,$$

where

$$\Sigma_w:=\mathrm{Hom}_{F_v}(K_w,F_v^{\mathrm{sep}})=\mathrm{Hom}_{F_v}(K_w,\mathfrak{F})=\{\sigma_{w,1},\dots,\sigma_{w,k_w}\}.$$

Hence,

$$\mathcal{C}(K,A,\mathfrak{F})=\{\ f:\coprod_{w|v}\Sigma_w\longrightarrow\mathbb{N}\mid\ \sum_{w|v}\sum_{i=1}^{k_w}f(\sigma_{w,i})=m\ \}.$$

The subgroup G_v of G also acts on the set $\mathcal{C}(K,A,\mathfrak{F})$.

Lemma 6.3.

(1) *The G_v-invariant subset $\mathcal{C}(K,A,\mathfrak{F})^{G_v}$ consists of functions f in $\mathcal{C}(K,A,\mathfrak{F})$ which are constant on Σ_w for all $w|v$.*

(2) *There is a commutative diagram*

$$\begin{tikzcd}
\mathcal{O}^*_{K_v,A_v} \arrow[r, "\otimes \mathrm{id}_{\mathfrak{F}}"] \arrow[d, "\psi_v"'] & \mathcal{O}^*_{K\otimes\mathfrak{F},A\otimes\mathfrak{F}} \arrow[d, "\phi"'] \\
\mathcal{C}(K,A,\mathfrak{F})^{G_v} \arrow[r] & \mathcal{C}(K,A,\mathfrak{F}).
\end{tikzcd}$$

Proof. The proof is the same as that of Lemma 6.2 and hence is omitted. ∎

Note that in Lemma 6.3, the map ψ_v is injective but it may not be surjective. To see this, we describe the map ψ_v. Let $\varepsilon_v : K_v \hookrightarrow A_v$ be an F_v-embedding. Let $A_v = \mathrm{Mat}_{ns_v}(D_v)$, where D_v is a central division F_v-algebra, and let $d_v := \deg(D_v)$. The map ε_v gives rise to an F_v-embedding $\varepsilon_w : K_w \hookrightarrow \mathrm{Mat}_{n_w}(D_v)$ for each place $w|v$ of K, where n_w are positive integers with $\sum_{w|v} n_w = ns_v$. Then the function $\psi_v([\varepsilon_v])$ is given by

$$\psi_v([\varepsilon_v])(\sigma)=n_w d_v/k_w, \quad \text{if } \sigma\in\Sigma_w.$$

From this, we see that the map ψ_v is injective. However, if f is a G_v-invariant function, say $f(\sigma)=a_w\in\mathbb{N}$ for $\sigma\in\Sigma_w$. Then the number $n_w := a_w k_w/d_w$ may not be an integer. This shows that the map ψ_v may not be surjective.

Lemma 6.4. *Let E/F be a finite Galois extension of global fields with Galois group H. For each v of F, let $H_v = \mathrm{Gal}(E_w/F_v)$ be a decomposition subgroup, where w is a place of K over v. Then $H = \langle H_v\rangle_{v\in V^F}$.*

Proof. This is a well-known consequence of the Chebotarev density theorem; we include the proof for the sake of completeness. Put $H_1 := \langle H_v\rangle_{v\in V^F}$. For any element $g\in H$, the Chebotarev density theorem tells us that there are infinitely many places v of F such that the Frobenius conjugacy class $Frob_v$ of v is equal to the conjugacy class C_g of g. This shows that the subgroup H_1 meets every conjugacy class of H as the subgroup H_v already intersects with the conjugacy class $Frob_v$. The lemma then follows from the next lemma, which is a basic exercise in group theory. ∎

Lemma 6.5. *Let H be a finite group and H_1 be a subgroup of H. Suppose that H_1 meets every conjugacy class of H. Then $H_1 = H$.*

Proof. Let $C \neq \{e\}$ be a conjugacy class of H. We want to bound its cardinality $|C|$. Let the group H act on itself by conjugation. Choose coset representatives $h_1,\dots h_r \in H$ for H/H_1, where $r=[H:H_1]$. Since $C\cap H_1\neq\emptyset$, let $x_0\in$

$C \cap H_1$ be an element. Let $O_H(x_0)$ and $O_{H_1}(x_0)$ be the H-orbit and H_1-orbit of x_0, respectively. We have

$$C = O_H(x_0), \quad \text{and} \quad C = \bigcup_{i=1}^{r} h_i O_H(x_0).$$

Then

$$|C| \le \sum_{i=1}^{r} |O_{H_1}(x_0)| \le [H : H_1] \cdot |C \cap H_1|,$$

as $O_{H_1}(x_0) \subset C \cap H_1$. It follows that

$$|H| - 1 = \sum_{C \ne \{e\}} |C| \le [H : H_1] \cdot (|H_1| - 1),$$

and hence that $H = H_1$. ∎

Proposition 6.6. *Let f be a function in $\mathcal{C}(K, A, \mathfrak{F})$. If $f \in \mathcal{C}(K, A, \mathfrak{F})^{G_v}$ for all $v \in V^F$, then $f \in \mathcal{C}(K, A, \mathfrak{F})^G$.*

PROOF. Let f be as in the proposition. The action of G factors through the finite quotient $H := \mathrm{Gal}(E/F)$, where E is the Galois closure of K over F in F^{sep}. Let $H_v := \mathrm{Gal}(EF_v/F_v) \subset H$ be the decomposition subgroup of v. Therefore, f is H_v-invariant for all $v \in V^F$. By Lemma 6.4 the group H is generated by H_v for all $v \in V^F$. Therefore, f is H-invariant. This completes the proof of the proposition. ∎

Now we are ready to prove our last main result.

Theorem 6.7. *Let $\iota_v : K_v \hookrightarrow A_v$ be an F_v-embedding of K_v into A_v for each $v \in V^F$. Suppose that $\iota_{v_1} \sim \iota_{v_2}$ for any two places $v_1, v_2 \in V^F$. Then there exist an F-embedding $\iota : K \hookrightarrow A$ such that $\iota \otimes \mathrm{id}_{F_v} \sim \iota_v$ (over F_v) for all $v \in V^F$.*

PROOF. Let ι_v be as given in the theorem. Let $f_v := \psi_v([\iota_v])$; it is a G_v-invariant function in $\mathcal{C}(K, A, \mathfrak{F})$. Since $[\iota_{v_1} \otimes \mathrm{id}_{\mathfrak{F}}] = [\iota_{v_1} \otimes \mathrm{id}_{\mathfrak{F}}] \in \mathcal{O}^*_{K\otimes\mathfrak{F}, A\otimes\mathfrak{F}}$, we have $f_{v_1} = f_{v_2}$ for any two places $v_1, v_2 \in V^F$. By Proposition 6.6, the function f_v is G-invariant and hence it is a constant function with value m/k. The map ι_v gives rise to maps $\iota_w : K_w \hookrightarrow \mathrm{Mat}_{n_w}(D_v)$ for some positive integers n_w with $\sum_{w|v} n_w = ns_v$. We get $f_v(\sigma) = n_w d_v / k_w = m/k = ns_v d_v / k$ and hence $n_w = ns_v k_w / k$ and $n_w c_w / k_w = ns_v c_w / k$. On the other hand, since we have an F_v-embedding $K_w \hookrightarrow \mathrm{Mat}_{n_W}(D_v)$, the condition $k_w | n_w c_w$ holds (Lemma 3.1). This shows that

$$x_w = \frac{ns_v c_w}{k} = \frac{n_w c_w}{k_w} \in \mathbb{N}, \quad \forall\, w \in V^K.$$

By Theorem 3.6, there is an F-embedding $\iota : K \hookrightarrow A$. Since the images $\psi_v([\iota \otimes \mathrm{id}_{F_v}])$ and $\psi_v([\iota_v])$ are the unique constant function, and the map ψ_v is injective, we get $[\iota \otimes \mathrm{id}_{F_v}] = [\iota_v]$, that is $\iota \otimes \mathrm{id}_{F_v} \sim \iota_v$ for all $v \in V^F$. This completes the proof of the theorem. ∎

Acknowledgments

The manuscript was revised during Yu's stay at the IEM, Universität Duisburg-Essen. He wishes to thank the IEM for kind hospitality and excellent working conditions. Yu was partially supported by grants NSC 97-2115-M-001-015-MY3, 100-2628-M-001-006-MY4 and AS-99-CDA-M01.

References

[1] 1967 Algebraic Number Theory. Edited by W.-S. Cassels and A. Fröhlich. (Proc. Instructional Conf., Brighton, 1965). Academic Press, 366 pp.

[2] P. Chuard-Koulmann and J. Morales, Extending involutions on Frobenius algebras. *Manuscripta Math.* **108** (2002), 439–451.

[3] M. Eichler, Zur Zahlentheorie der Quaternionen-Algebren. *J. Reine Angew. Math.* **195** (1955), 127–151.

[4] R. S. Pierce, *Associative algebras.* Graduate Texts in Mathematics, **88**. Springer-Verlag, New York-Berlin, 1982. 436 pp.

[5] G. Prasad and A. Rapinchuk, Computation of the metaplectic kernel. *Inst. Hautes Études Sci. Publ. Math.* **84** (1996), 91–187.

[6] G. Prasad and A. Rapinchuk, Local-global principles for embedding of fields with involution into simple algebras with involution, *Comment. Math. Helv.* **85** (2010), 583–645.

[7] I. Reiner, *Maximal orders.* London Mathematical Society Monographs, No. **5**. Academic Press, London-New York, 1975. 395 pp.

[8] J.-P. Serre, *Local fields.* **GTM 67**, Springer-Verlag, 1979.

[9] M.-F. Vignéras, Arithmétique des algèbres de quaternions. Lecture Notes in Math., vol. 800, Springer-Verlag, 1980.

[10] C.-F. Yu, Construction of Galois covers of curves with groups of SL_2-type. *C. R. Acad. Sci. Paris Sér. I Math.* **345** (2007), 77-80.

[11] C.-F. Yu, Embeddings of fields into simple algebras: generalizations and applications. arXiv:1105.2895. 18 pp.

(Shih) Institute of Mathematics, Academia Sinica, Astronomy Mathematics Building, No. 1, Roosevelt Rd. Sec. 4, Taipei, Taiwan, 10617
E-mail address: `r95221018@ntu.edu.tw`

(Yang) Department of Mathematics, National Taiwan University, Astronomy Mathematics Building, No. 1, Roosevelt Rd. Sec. 4, Taipei, Taiwan, 10617
E-mail address: `d97221007@ntu.edu.tw`

(Yu) Institute of Mathematics, Academia Sinica and NCTS (Taipei Office), Astronomy Mathematics Building, No. 1, Roosevelt Rd. Sec. 4, Taipei, Taiwan, 10617
E-mail address: `chiafu@math.sinica.edu.tw`